

***The Yucca Mountain Dump Plan
Would Launch Up to 341 Barges of
Deadly High-Level Radioactive Waste
Onto the Waters of Florida's Atlantic Coastline***

As part of its plan to transport high-level radioactive waste to Western Shoshone Indian land at Yucca Mountain, Nevada, the U.S. Department of Energy (DOE) proposes up to 341 barges carrying giant high-level radioactive waste containers onto the Atlantic along Florida's coastline, into Fort Lauderdale and Miami. See the second page of this fact sheet for a map of the proposed route and a breakdown of shipment numbers.

Accidents happen. But what if high-level radioactive waste is involved? U.S. Nuclear Regulatory Commission (NRC) design criteria for atomic waste transport containers are woefully inadequate. Rather than full-scale physical safety testing, scale model tests and computer simulations are all that is required.

The underwater immersion design criteria are meant to "test" (on paper, at least) the integrity of a slightly damaged container submerged under 3 feet of water for 8 hours. An undamaged cask is "tested" (on computers, at least) for a 1 hour submersion under 656 feet of water.

But if a cask were accidentally immersed under water, or sunk by terrorists, is it reasonable for NRC to assume that the cask would only be slightly damaged, or not damaged at all? Given that barge casks could weigh well over 100 tons (even up to 140 tons), how can NRC assume that they could be recovered from underwater within 1 hour, or even within 8 hours? Special cranes capable of lifting such heavy loads would have to be located, brought in, and set up. And what about submersions that occur at depths deeper than 656 feet underwater?

The dangers of nuclear waste cask submersion underwater are two fold. First, radioactivity could leak from the cask into the water. Each container would hold 200 times the long-lasting radioactivity released by the Hiroshima atomic bomb. Given high-level atomic waste's deadliness, leakage of even a fraction of a cask's contents could spell unprecedented catastrophe for a vast stretch of the Florida coastline. Second, enough fissile uranium-235 and plutonium is present in high-level atomic waste that water, with its neutron moderating properties, could actually cause a nuclear chain reaction to take place within the cask. Such an inadvertent criticality event in Sept. 1999 at a nuclear fuel factory in Japan led to the deaths of two workers; many hundreds of nearby residents, including children, received radiation doses well above safety standards.

STOP THE ACCIDENT BEFORE IT HAPPENS!


***Don't let D.O.E. and N.R.C. get away with
shipping high-level radioactive wastes on the California coastline!***

**Urge Your U.S. Senators and Representative to oppose
the Yucca Mountain dump plan!**

Call their offices via the U.S. Capitol Switchboard: 202.224.3121.

**For more information, contact Nuclear Information & Resource Service, 301.270.6477,
nirsnet@nirs.org, www.nirs.org**

Barge Shipments of High-Level Radioactive Waste on Florida's Atlantic Coast
 Proposed by U.S. Dept. of Energy under its Yucca Mountain Plan



Map taken from Figure J-9, Routes analyzed for barge transportation from sites to nearby railheads, page J-79.

<u>Nuclear Reactor</u>	<u>Location</u>	<u>Shipment #'s Proposed</u>	<u>Barges offloaded at:</u>
St. Lucie 1	Hutchinson Island	Up to 16	Port Everglades, Ft. Lauderdale
St. Lucie 2	Hutchinson Island	Up to 150	Port Everglades, Ft. Lauderdale
Turkey Point 3	Florida City	Up to 87	Port of Miami
Turkey Point 4	Florida City	Up to 88	Port of Miami
Totals		Up to 341	

Table taken from Table J-27, Barge shipments and ports, page J-83.

Map and table taken from U.S. Department of Energy, "Final Environmental Impact Statement for Yucca Mountain," Appendix J ("Transportation"), Feb. 2002.