

***The Yucca Mountain Dump Plan
Would Launch Up to 326 Barges of
Deadly High-Level Radioactive Waste
Onto the Waters of the Chesapeake Bay***

As part of its plan to transport high-level radioactive waste to Western Shoshone Indian land at Yucca Mountain, Nevada, the U.S. Department of Energy (DOE) proposes up to 326 barges carrying giant high-level radioactive waste containers onto the waters of the Chesapeake Bay from the Calvert Cliffs nuclear power plant to the Port of Baltimore. See the second page of this fact sheet for a map of the proposed route.

Accidents happen. But what if high-level radioactive waste is involved? U.S. Nuclear Regulatory Commission (NRC) design criteria for atomic waste transport containers are woefully inadequate. Rather than full-scale physical safety testing, scale model tests and computer simulations are all that is required.

The underwater immersion design criteria are meant to “test” (on paper, at least) the integrity of a slightly damaged container submerged under 3 feet of water for 8 hours. An undamaged cask is “tested” (on computers, at least) for a 1 hour submersion under 656 feet of water.

But if a cask were accidentally immersed under water, or sunk by terrorists, is it reasonable for NRC to assume that the cask would only be slightly damaged, or not damaged at all? Given that barge casks could weigh well over 100 tons (even up to 140 tons), how can NRC assume that they could be recovered from underwater within 1 hour, or even within 8 hours? Special cranes capable of lifting such heavy loads would have to be located, brought in, and set up.

The dangers of nuclear waste cask submersion underwater are two fold. First, radioactivity could leak from the cask into the water. Each barge sized container could hold 200 times the long-lasting radioactivity given by the Hiroshima atomic bomb. Given high-level atomic waste’s deadliness, leakage of even a fraction of a cask’s contents into the Chesapeake Bay could spell unprecedented catastrophe. Second, enough fissile uranium-235 and plutonium is present in high-level atomic waste that water, with its neutron moderating properties, could actually cause a nuclear chain reaction to take place within the cask. Such an inadvertent criticality event in Sept. 1999 at a nuclear fuel factory in Japan led to the deaths of two workers; many hundreds of nearby residents, including children, received radiation doses well above safety standards.

STOP THE ACCIDENT BEFORE IT HAPPENS!


***Don’t let D.O.E. and N.R.C. get away with
shipping high-level radioactive wastes on the Chesapeake Bay!***

**Urge Your U.S. Senators and Representative to oppose
the Yucca Mountain dump plan!**

Call their offices via the U.S. Capitol Switchboard: 202.224.3121.

**For more information, contact Nuclear Information & Resource Service, 301.270.6477,
nirsnet@nirs.org, www.nirs.org**

Barge Shipments of High-Level Radioactive Waste on the Chesapeake Bay Proposed by U.S. Dept. of Energy under its Yucca Mountain Plan



Map taken from Figure J-9, Routes analyzed for barge transportation from sites to nearby railheads, page J-78.

<u>Nuclear Reactor</u>	<u>Location</u>	<u>Number of Shipments Proposed</u>	<u>Barges offloaded at:</u>
Calvert Cliffs 1	Lusby, MD	323	Port of Baltimore
Calvert Cliffs 2	Lusby, MD	3	Port of Baltimore
Total		Up to 326	

Table taken from Table J-27, Barge shipments and ports, page J-83.

Map and table taken from U.S. Department of Energy, "Final Environmental Impact Statement for Yucca Mountain," Appendix J ("Transportation"), Feb. 2002.